



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.


OZ , and $\beta \cos\phi$ about OX . The former does not alter the position of OC , while the latter turns OC in a time t through an angle $\beta \cos\phi t$. Hence the body is acted upon by a westerly component, due to the change of direction of gravity, $=g \sin(\beta \cos\phi t) = g \beta \cos\phi t$, since βt is small. Now let y be the distance the body is in space from the plane XZ at the moment the body begins to fall, and then the equation of motion of the body in space is $\frac{d^2y}{dt^2} = g \beta t \cos\phi$. Integrating this, and remembering that, as explained above, $\frac{dy}{dt} = -\beta H \cos\phi$, we get

$y = -\beta H t \cos\phi + \frac{1}{6} g \beta t^3 \cos\phi$. $\therefore E_d = \beta t \cos\phi (H - \frac{1}{3} g t^2)$. But $\frac{1}{2} g t^2 = H + \Delta$ and the centrifugal force at the equator $= \beta^2 r = \frac{4\pi^2 r}{T^2}$, where r = radius of the earth.

$$\therefore \beta^2 = \frac{4\pi^2}{T^2} \text{ and } \beta = \frac{2\pi}{T}. \text{ Substituting we get}$$

$$E_d = \frac{2\pi t \cos\phi}{T} \{ H - \frac{1}{3}(H + \Delta) \} = \frac{2\pi t \cos\phi}{T} (\frac{2}{3}H - \frac{1}{3}\Delta)$$

$$= \frac{2\pi t \cos\phi}{3T} (2H - \Delta) = \frac{4\pi t (H - \frac{1}{3}\Delta) \cos\phi}{3T}.$$



[NOTE.—No solution has yet been received to problem 17. EDITOR.]

PROBLEMS.

24. Proposed by Professor J. F. W. SCHEFFER, A. M., Hagerstown, Maryland.

A sphere whose center of gravity does not coincide with its geometrical center is placed on a rough inclined plane. State under what circumstances the sphere will slide without rolling, roll without sliding, and neither roll nor slide.

25. Proposed by Professor GEORGE LILLEY, LL. D., Ex-President of Washington State Agricultural College and School of Science, Portland, Oregon.

It is known that if the velocity of a certain freight train is 30 miles an hour it can be brought to a stand still in a distance of 500 feet by setting the brakes. It was stopped in 1200 feet by setting the brakes. Find its velocity, the forces exerted by the brakes being the same in each case.

